

References for *Antioxidants in food, drinks and supplements for cardiovascular health*

1. National Heart Foundation of Australia. Summary of evidence. Antioxidants in food, drinks and supplements for cardiovascular health. Melbourne: National Heart Foundation of Australia, 2009.
2. National Heart Foundation of Australia. Position statement. The use of antioxidant supplements in heart disease. Melbourne: National Heart Foundation of Australia, 2002.
3. National Health and Medical Research Council. Australian Guidelines to Reduce Health Risks From Drinking Alcohol. Canberra: National Health and Medical Research Council, 2009.
4. National Health and Medical Research Council. Nutrient Reference Values for Australia and New Zealand including Recommended Dietary Intakes. Canberra: National Health and Medical Research Council, 2006. Page 248.
5. National Heart Foundation of Australia. Reducing risk in heart disease. Guidelines for preventing cardiovascular events in people with coronary heart disease. Melbourne: National Heart Foundation of Australia, 2008.
6. National Heart Foundation of Australia. Guidelines for the prevention, detection and management of chronic heart failure in Australia. Melbourne: National Heart Foundation of Australia, 2006.
7. SIGNAL and National Public Health Partnership. An Intervention Portfolio To Promote Fruit and Vegetable Consumption. Part 1 – The Process and Portfolio, 2001.
8. Preventative Health Taskforce – Obesity Working Group: National Preventative Health Strategy. Obesity in Australia: a need for urgent action. Canberra: Preventative Health Taskforce, 2009.
9. Pollard CM, Lewis JM, Binns CW. Selecting interventions to promote fruit and vegetable consumption: from policy to action, a planning framework case study in Western Australia. Aust New Zealand Health Policy 2008; 5: 27.
10. National Health and Medical Research Council. A guide to the development, implementation and evaluation of clinical practice guidelines. Canberra: National Health and Medical Research Council, 1999.
11. Appel L, Moore T, Obarzanek E et al. A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research Group. N Engl J Med 1997; 336: 1117–24.

12. Most MM. Estimated phytochemical content of the dietary approaches to stop hypertension (DASH) diet is higher than in the Control Study Diet. *J Am Diet Assoc* 2004; 104: 1725–7.
13. National Heart Lung and Blood Institute. Your Guide to Lowering Your Blood Pressure with DASH. NIH Publication No. 06–5834 NIH, 2006.
14. Rissanen TH, Voutilainen S, Virtanen JK et al. Low intake of fruits, berries and vegetables is associated with excess mortality in men: the Kuopio Ischaemic Heart Disease Risk Factor (KIHD) Study. *J Nutr* 2003; 133: 199–204.
15. Nakamura K, Nagata C, Oba S et al. Fruit and vegetable intake and mortality from cardiovascular disease are inversely associated in Japanese women but not in men. *J Nutr* 2008; 138: 1129–34.
16. He FJ, Nowson CA, MacGregor GA. Fruit and vegetable consumption and stroke: meta-analysis of cohort studies. *Lancet* 2006; 367: 320–6.
17. Law MR, Morris JK. By how much does fruit and vegetable consumption reduce the risk of ischaemic heart disease? *Eur J Clin Nutr* 1998; 52: 549–56.
18. Knekt P, Jarvinen R, Reunanen A et al. Flavonoid intake and coronary mortality in Finland: a cohort study. *BMJ* 1996; 312: 478–81.
19. Mink PJ, Scrafford CG, Barraj LM et al. Flavonoid intake and cardiovascular disease mortality: a prospective study in postmenopausal women. *Am J Clin Nutr* 2007; 85: 895–909.
20. He FJ, Nowson CA, Lucas M et al. Increased consumption of fruit and vegetables is related to a reduced risk of coronary heart disease: meta-analysis of cohort studies. *J Hum Hypertens* 2007; 21: 717–28.
21. Dauchet L, Amouyel P, Hercberg S et al. Fruit and vegetable consumption and risk of coronary heart disease: a meta-analysis of cohort studies. *J Nutr* 2006; 136: 2588–93.
22. Joshipura KJ, Hu FB, Manson JE et al. The effect of fruit and vegetable intake on risk for coronary heart disease. *Ann Intern Med* 2001; 134: 1106–14.
23. Hertog MG, Feskens EJ, Hollman PC et al. Dietary antioxidant flavonoids and risk of coronary heart disease: the Zutphen Elderly Study. *Lancet* 1993; 342: 1007–11.
24. Arts IC, Jacobs DR Jr, Harnack LJ et al. Dietary catechins in relation to coronary heart disease death among postmenopausal women. *Epidemiology* 2001; 12: 668–75.
25. Alexopoulos N, Vlachopoulos C, Aznaouridis K et al. The acute effect of green tea consumption on endothelial function in healthy individuals. *Eur J Cardiovasc Prev Rehabil* 2008; 15: 300–5.

26. Nagaya N, Yamamoto H, Uematsu M et al. Green tea reverses endothelial dysfunction in healthy smokers. *Heart* 2004; 90: 1485–6.
27. Kim W, Jeong MH, Cho SH et al. Effect of green tea consumption on endothelial function and circulating endothelial progenitor cells in chronic smokers. *Circ J* 2006; 70: 1052–7.
28. Jochmann N, Lorenz M, Krosigk A et al. The efficacy of black tea in ameliorating endothelial function is equivalent to that of green tea. *Br J Nutr* 2008; 99: 863–8.
29. Hodgson JM, Croft KD, Mori TA et al. Regular ingestion of tea does not inhibit in vivo lipid peroxidation in humans. *J Nutr* 2002; 132: 55–8.
30. Duffy SJ, Keaney JF Jr, Holbrook M et al. Short- and long-term black tea consumption reverses endothelial dysfunction in patients with coronary artery disease. *Circulation* 2001; 104: 151–6.
31. Hodgson JM, Burke V, Puddey IB. Acute effects of tea on fasting and postprandial vascular function and blood pressure in humans. *J Hypertens* 2005; 23: 47–54.
32. Grassi D, Mulder T, Draijer R et al. Black tea consumption dose-dependently improves flow-mediated dilation in healthy males. *J Hypertens* 2009; 27: 774–81.
33. Widlansky ME, Hamburg NM, Anter E et al. Acute EGCG supplementation reverses endothelial dysfunction in patients with coronary artery disease. *J Am Coll Nutr* 2007; 26: 95–102.
34. Loke WM, Hodgson JM, Proudfoot JM et al. Pure dietary flavonoids quercetin and (-)-epicatechin augment nitric oxide products and reduce endothelin-1 acutely in healthy men. *Am J Clin Nutr* 2008; 88: 1018–25.
35. Nagao T, Komine Y, Soga S et al. Ingestion of a tea rich in catechins leads to a reduction in body fat and malondialdehyde-modified LDL in men. *Am J Clin Nutr* 2005; 81: 122–9.
36. Kajimoto O, Kajimoto Y, Yabune M et al. Tea catechins with a galloyl moiety reduce body weight and fat. *J Health Sci* 2005; 1: 161–71.
37. Nagao T, Hase T, Tokimitsu I. A green tea extract high in catechins reduces body fat and cardiovascular risks in humans. *Obesity (Silver Spring)* 2007; 15: 1473–83.
38. Maki KC, Reeves MS, Farmer M et al. Green tea catechin consumption enhances exercise-induced abdominal fat loss in overweight and obese adults. *J Nutr* 2009; 139: 264–70.
39. Diepvens K, Westerterp KR, Westerterp-Plantenga MS. Obesity and thermogenesis related to the consumption of caffeine, ephedrine, capsaicin, and green tea. *Am J Physiol Regul Integr Comp Physiol* 2007; 292: R77–85.

40. Peters U, Poole C, Arab L. Does Tea Affect Cardiovascular Disease? A Meta-Analysis. *Am J Epidemiol* 2001; 154: 495–503.
41. Stangl V, Lorenz M, Stangl K. The role of tea and tea flavonoids in cardiovascular health. *Mol Nutr Food Res* 206; 50: 218–28.
42. Khan N, Mukhtar H. Tea polyphenols for health promotion. *Life Sci* 2007; 81: 519–33.
43. Kuriyama S. The relation between green tea consumption and cardiovascular disease as evidenced by epidemiological studies. *J Nutr* 2008; 138: 1548S–1553S.
44. Wolfram S. Effects of green tea and EGCG on cardiovascular and metabolic health. *J Am Coll Nutr* 2007; 26: 373S–388S.
45. Huxley RR, Neil HA. The relation between dietary flavonol intake and coronary heart disease mortality: a meta-analysis of prospective cohort studies. *Eur J Clin Nutr* 2003; 57: 904–8.
46. Quinlan P, Lane J, Aspinall L. Effects of hot tea, coffee and water ingestion on physiological responses and mood: the role of caffeine, water and beverage type. *Psychopharmacology (Berl)* 1997; 134: 164–73.
47. Hodgson JM. Effects of tea and tea flavonoids on endothelial function and blood pressure: a brief review. *Clin Exp Pharmacol Physiol* 2006; 33: 838–41.
48. Stensvold I, Tverdal A, Solvoll K et al. Tea consumption. Relationship to cholesterol, blood pressure, and coronary and total mortality. *Prev Med* 1992; 21: 546–53.
49. Hakim IA, Alsaif MA, Alduwaihy M et al. Tea consumption and the prevalence of coronary heart disease in Saudi adults: results from a Saudi national study. *Prev Med* 2003; 36: 64–70.
50. Yang YC, Lu FH, Wu JS et al. The protective effect of habitual tea consumption on hypertension. *Arch Intern Med* 2004; 164: 1534–40.
51. Hodgson JM, Puddey IB, Burke V et al. Effects on blood pressure of drinking green and black tea. *J Hypertens* 1999; 17: 457–63.
52. Bingham SA, Vorster H, Jerling JC et al. Effect of black tea drinking on blood lipids, blood pressure and aspects of bowel habit. *Br J Nutr* 1997; 78: 41–55.
53. Taubert D, Roesen R, Schomig E. Effect of cocoa and tea intake on blood pressure: a meta-analysis. *Arch Intern Med* 2007; 167: 626–34.
54. Hooper L, Kroon PA, Rimm EB et al. Flavonoids, flavonoid-rich foods, and cardiovascular risk: a meta-analysis of randomized controlled trials. *Am J Clin Nutr* 2008; 88: 38–50.
55. Corti R, Flammer AJ, Hollenberg NK et al. Cocoa and cardiovascular health. *Circulation* 2009; 119: 1433–41.

56. Buijsse B, Feskens EJ, Kok FJ et al. Cocoa intake, blood pressure, and cardiovascular mortality: the Zutphen Elderly Study. *Arch Intern Med* 2006; 166: 411–7.
57. Taubert D, Roesen R, Lehmann C et al. Effects of low habitual cocoa intake on blood pressure and bioactive nitric oxide: a randomized controlled trial. *JAMA* 2007; 298: 49–60.
58. Kris-Etherton PM, Keen CL. Evidence that the antioxidant flavonoids in tea and cocoa are beneficial for cardiovascular health. *Curr Opin Lipidol* 2002; 13: 41–9.
59. Steinberg FM, Bearden MM, Keen CL. Cocoa and chocolate flavonoids: implications for cardiovascular health. *J Am Diet Assoc* 2003; 103: 215–23.
60. Engler MB, Engler MM. The emerging role of flavonoid-rich cocoa and chocolate in cardiovascular health and disease. *Nutr Rev* 2006; 64: 109–18.
61. Mehrinfar R, Frishman WH. Flavanol-rich cocoa: a cardioprotective nutraceutical. *Cardiol Rev* 2008; 16: 109–15.
62. Keen CL, Holt RR, Oteiza PI et al. Cocoa antioxidants and cardiovascular health. *Am J Clin Nutr* 2005; 81: 298S–303.
63. Jee SH, He J, Whelton PK et al. The effect of chronic coffee drinking on blood pressure: a meta-analysis of controlled clinical trials. *Hypertension* 1999; 33: 647–52.
64. Noordzij M, Uiterwaal CS, Arends LR et al. Blood pressure response to chronic intake of coffee and caffeine: a meta-analysis of randomized controlled trials. *J Hypertens* 2005; 23: 921–8.
65. Weusten-Van der Wouw MP, Katan MB, Viani R et al. Identity of the cholesterol-raising factor from boiled coffee and its effects on liver function enzymes. *J Lipid Res* 1994; 35: 721–33.
66. Heckers H, Gobel U, Kleppel U. End of the coffee mystery: diterpene alcohols raise serum low-density lipoprotein cholesterol and triglyceride levels. *J Intern Med* 1994; 235: 192–3.
67. Urgert R, van der Weg G, Kosmeijer-Schuil TG et al. Levels of the Cholesterol-Elevating Diterpenes Cafestol and Kahweol in Various Coffee Brews. *Journal of Agricultural and Food Chemistry* 2002; 43: 2167–2172.
68. Jee SH, He J, Appel LJ et al. Coffee consumption and serum lipids: a meta-analysis of randomized controlled clinical trials. *Am J Epidemiol* 2001; 153: 353–62.
69. Sofi F, Conti AA, Gori AM et al. Coffee consumption and risk of coronary heart disease: a meta-analysis. *Nutr Metab Cardiovasc Dis* 2007; 17: 209–23.

70. Myers MG, Basinski A. Coffee and coronary heart disease. *Arch Intern Med* 1992; 152: 1767–72.
71. Greenland S. A meta-analysis of coffee, myocardial infarction, and coronary death. *Epidemiology* 1993; 4: 366–74.
72. Kawachi I, Colditz GA, Stone CB. Does coffee drinking increase the risk of coronary heart disease? Results from a meta-analysis. *Br Heart J* 1994; 72: 269–75.
73. Wu JN, Ho SC, Zhou C et al. Coffee consumption and risk of coronary heart diseases: A meta-analysis of 21 prospective cohort studies. *Int J Cardiol* 2009; 137: 216–25.
74. Mukamal KJ, Hallqvist J, Hammar N et al. Coffee consumption and mortality after acute myocardial infarction: The Stockholm Heart Epidemiology Program. *American Heart Journal* 2009; 157: 495–501.
75. Ahmed HN, Levitan EB, Wolk A et al. Coffee consumption and risk of heart failure in men: An analysis from the Cohort of Swedish Men. *American Heart Journal* 2009; In press, corrected proof.
76. Lopez-Garcia E, van Dam RM, Li TY et al. The relationship of coffee consumption with mortality. *Ann Intern Med* 2008; 148: 904–14.
77. Geleijnse JM. Habitual coffee consumption and blood pressure: an epidemiological perspective. *Vasc Health Risk Manag* 2008; 4: 963–70.
78. van Dam RM, Hu FB. Coffee consumption and risk of type 2 diabetes: a systematic review. *JAMA* 2005; 294: 97–104.
79. Katsiki N, Manes C. Is there a role for supplemented antioxidants in the prevention of atherosclerosis? *Clin Nutr* 2009; 28: 3–9.
80. Shekelle PG, Morton SC, Jungvig LK et al. Effect of supplemental vitamin E for the prevention and treatment of cardiovascular disease. *J Gen Intern Med* 2004; 19: 380–9.
81. Vivekananthan DP, Penn MS, Sapp SK et al. Use of antioxidant vitamins for the prevention of cardiovascular disease: meta-analysis of randomised trials. *Lancet* 2003; 361: 2017–23.
82. Eidelman RS, Hollar D, Hebert PR et al. Randomized trials of vitamin E in the treatment and prevention of cardiovascular disease. *Arch Intern Med* 2004; 164: 1552–6.
83. Ye Z, Song H et al. Antioxidant vitamins intake and the risk of coronary heart disease: meta-analysis of cohort studies. *Eur J Cardiovasc Prev Rehabil* 2008; 15: 26–34.
84. Knekt P, Ritz J, Pereira MA et al. Antioxidant vitamins and coronary heart disease risk: a pooled analysis of 9 cohorts. *Am J Clin Nutr* 2004; 80: 1508–20.