

Heart information



Heart valve surgery



Contents

Page 1	What are heart valves?
Page 3	What is heart valve surgery?
Page 3	Valve repair
Page 4	Valve replacement
Page 4	What happens during heart valve surgery?
Page 5	Why do I need heart valve surgery?
Page 6	What will happen after heart valve surgery?
Page 6	Will I feel pain after the operation?
Page 6	How long will I have to stay in hospital?
Page 7	How long will it take to recover from heart valve surgery?
Page 7	When can I start eating again?
Page 8	What should I be eating?
Page 8	What if I am constipated?
Page 8	How active can I be?
Page 9	When can I be more physically active?
Page 9	When can I have sex?
Page 10	Why am I so tired?
Page 10	Why do I feel great one day, but dreadful the next?
Page 10	When can I return to work?
Page 10	Palpitations
Page 11	Heart attack
Page 12	Endocarditis
Page 12	Will I need to take medicine?
Page 13	Important things to remember about anti-coagulant medicine
Page 14	Warfarin and food
Page 15	What should I talk to my doctor about?
Page 16	Glossary

What are heart valves?

Heart valves are like one-way doors that control the direction of blood flow between the four chambers of the heart. There are two upper chambers (the atria) and two lower chambers (the ventricles) (see the diagram on page 2). The atria receive blood from the body and pump it into the ventricles. The ventricles pump blood away from the heart to the rest of the body.

With each heartbeat, blood is squeezed from the atria into the ventricles, and then out of the ventricles to the rest of the body. After each contraction of the atria or the ventricles, the inner flaps of the valves (often called 'cusps' or 'leaflets') come together to seal off the chambers of the heart and make sure that the blood does not flow backwards. They are forced open again by the normal pumping action of the heart to allow forward blood flow.


The four heart valves are:

- the **tricuspid valve**, on the right side of the heart, which controls the blood flow between the right atrium and the right ventricle
- the **mitral valve**, on the left side of the heart, which controls the blood flow between the left atrium and the left ventricle
- the **pulmonary valve**, on the right side of the heart, which separates the right ventricle and the pulmonary artery – the blood vessel that carries blood from the heart to the lungs
- the **aortic valve**, on the left side of the heart, which separates the left ventricle and the aorta.

For more
information
on this topic
please call
1300 36 27 87



The diagram below shows the four chambers of the heart and the location of the valves. It also shows, using arrows, how blood flows through the heart.



Blood that has circulated through your body has had most of its oxygen and nutrients removed. It returns into the right atrium, passes through the tricuspid valve into the right ventricle, and is then pumped through the pulmonary valve into the vessels leading to your lungs.

The lungs add needed oxygen to the blood and remove carbon dioxide waste. Now the blood returns to your heart, flowing from the lungs into the left atrium. From the left atrium, the blood passes through the mitral valve into the left ventricle. Finally, it's pumped through the aortic valve into blood vessels supplying the rest of your body.

What is heart valve surgery?

Heart valve surgery is an operation to fix a damaged or faulty heart valve. It usually takes between four and six hours.

There are two main types of heart valve surgery that you can have: valve repair and valve replacement.

Valve repair

Sometimes a faulty valve can be repaired by cutting away excess tissue in the cusps of the valve and sewing the edges together. It can also be repaired by shortening or connecting the cords that act like hinges on the valve.

Special rings called 'prosthetic rings' or 'annuloplasty rings' (see diagram below) can also be used to narrow an enlarged valve and strengthen the repair.

Another method used to open a narrowed valve is called 'balloon valvotomy'. In this procedure, a small tube (a 'catheter') is inserted into an artery in your arm or groin and guided into your heart. Once it is positioned within the problem valve, a special balloon at the tip of the catheter is inflated to stretch open the valve. The catheter and balloon are then removed.

An advantage of valve repair operations is that usually your own valve tissues are used in the repair.



Valve replacement

If a valve can't be repaired, it is sometimes removed and replaced with a new valve. The new valve is sewn onto a rim of tissue that is kept from the original valve.

Several types of replacement valves are used. They fall into two groups.

Biological tissue valves

These are human or animal valves. Animal valves (usually taken from a cow or pig) are mounted in a cloth-covered metal or plastic frame to make them easier to insert. Human valves are taken from donated human hearts and can be sewn directly into place.

Mechanical valves

These have some advantages over tissue valves because they last longer. However, blood clots can form on the synthetic material. To overcome this, anti-clotting medicine (see page 12) needs to be taken for life.

What happens during heart valve surgery?

During the operation, your surgeon will cut down the midline of your chest, through your breastbone, to reach your heart. Your body will be kept cool to protect your vital organs by slowing down their working rate so that they need less oxygen. A heart-lung machine will take over the function of your heart and lungs.

A note about blood transfusions

If you object to having blood transfusions, please tell your doctor and surgeon before surgery.

If you need a blood transfusion, all blood products used for transfusion in Australia are strictly screened to protect patients against viruses that can cause hepatitis and AIDS. Sometimes you can arrange to donate some of your own blood before you have surgery, so that you can use it if you need a blood transfusion. However, this isn't always possible.

Why do I need heart valve surgery?

You need heart valve surgery because your heart valves are not working properly and your doctor has diagnosed you with heart valve damage.


Sometimes people are born with damaged valves, which may be fixed soon after birth or later in life.

Disease or infection can also damage your valves. Rheumatic fever is a frequent cause of valve damage. It makes a valve gradually become stiff so that it doesn't open and close properly.

A damaged valve that limits forward blood flow by not opening properly is called a 'stenotic valve' (see diagram below).

A damaged valve that doesn't close properly, allowing blood to flow back into the chamber, is called an 'incompetent valve' or a 'regurgitant valve'. Often both conditions are present.

When a valve is damaged, your heart cannot pump blood efficiently. It has to work harder to deliver oxygen-rich blood around your body. An overworked heart may begin to weaken, causing shortness of breath, pain, tiredness and a build-up of fluid in the body. If this happens, the valve may have to be replaced or repaired (see pages 3 and 4).



What will happen after heart valve surgery?

After the operation, you will be taken to a recovery area or intensive care unit for close supervision until you wake up from the anaesthetic. You will probably stay in the intensive care unit for one or two days. After the intensive care unit, you will be moved to another ward until you are ready to go home.

Before you leave hospital, your doctor will make follow-up appointments for you to discuss your medicines, wound care and activities with them. Your cardiologist will continue to oversee your general heart health. Your first outpatient appointment will usually be two to four weeks after you leave hospital.

About four to six weeks after leaving hospital, you will see your surgeon, or a member of the surgical team. Your doctor will check your wound and general progress, and you may also have an X-ray at this time.

Quick tip

Sitting up and moving your arms and shoulders frequently may help to reduce your soreness.

Will I feel pain after the operation?

You will probably feel quite sore, especially in the first few days after the operation. You will be given painkillers regularly to make sure that you don't feel severe pain. If the pain starts to build up, tell your nurse sooner rather than later. You and your nurse need to communicate and manage your pain together.

Medicine won't get rid of all of your pain, but it can manage it. Stronger painkilling medicines are needed for only a few days, so there is very little risk of becoming addicted to them.

How long will I have to stay in hospital?

This varies depending on your condition and your response to surgery. On average, after heart valve surgery, people stay in hospital for six to nine days.

How long will it take to recover from heart valve surgery?

Normal recovery from heart surgery takes four to six weeks. During this time, you will begin to strengthen your muscles and return to your usual activities.

Soon after your operation, your chest wound will be exposed to the air, allowing it to dry. After a few days, the wound can be cleaned.

Sometimes wire or special strips of tape are used to hold the breastbone together. These do not need to be removed. It takes about six to 12 weeks for the breastbone to heal completely. During that time you shouldn't lift anything heavy.

After major surgery, such as heart valve surgery, recovery time may seem to pass slowly. Your body has been slowed down by the lowered activity, lack of good sleep, medicines and the surgery itself. You may feel drained, physically and emotionally.

Some hospitals run outpatient rehabilitation programs. These can be a very important part of your recovery and will help you to continue the gradual increase in activity you began in hospital.

When can I start eating again?

After the operation, while you are still under the anaesthetic, a breathing tube is put into your windpipe. The tube is usually removed within eight to 24 hours, and most of that time you will have been asleep. After the tube is removed, you will be able to swallow a small amount of liquid and build up to foods over the next couple of days.



Cardiac rehabilitation

The Heart Foundation and World Health Organization recommend that all patients who have had a heart attack, heart surgery, coronary angioplasty, angina or other heart and blood vessel disease attend an appropriate cardiac rehabilitation and prevention program.



Want to know
more? Call
1300 36 27 87



What should I be eating?

Enjoying a variety of foods from the different food groups is the key to healthy eating. Choose:

- mainly plant-based foods, such as vegetables, fruit, legumes (e.g. chick peas, kidney beans, baked beans and lentils) and plain unsalted nuts
- wholegrain cereal foods, such as breakfast cereals, bread, pasta, noodles and rice
- moderate amounts of lean unprocessed meats, skinless poultry and reduced fat dairy products
- fish – include a 150 gram serve in two or three meals per week
- foods with low or reduced salt content and foods labelled ‘no added salt’, ‘low salt’ or ‘salt reduced’
- plant-based oils, such as canola, sunflower, soybean, olive and peanut oils.

What if I am constipated?

Unfortunately, many pain-relieving tablets can make you constipated. The best remedy is to eat high-fibre foods, such as bran, seeds, fruit and vegetables, and drink plenty of fluids (aim for 6–8 glasses per day). Follow your exercise program, because exercise will help promote regularity.

How active can I be?

This will depend on your recovery. You will probably start just by sitting in a chair or walking around the room. Later, there will be short walks in the corridor and, eventually, stair climbing and brisk longer walks to prepare for home. Sponge baths are given right away and within a few days, you will be able to take a shower and wash your hair.

When can I be more physically active?

Once you are at home, continue with the level of activities that you started in hospital. Aim to progress your activity gradually over the weeks and months ahead as advised by your surgical team.

Set realistic goals for yourself. Don't over do it, but don't be inactive either. Rest when you need to and change an activity if it is making you too tired. Doing too much at this time won't injure the heart. It will, however, make you very tired.

If you experience chest wound complications, such as redness or weeping and/or movement of the breastbone, you should contact your surgical team for advice. You may need to stop your activity or take it a bit more slowly and allow more time for healing.

As you become more active during your recovery, you will probably get some pain associated with the wound or muscles in your chest, shoulder, neck and back. Continue to take your pain-relieving tablets (e.g. paracetamol) when you need them. You will be able to reduce these gradually. **Don't neglect any type of severe pain.**

When can I have sex?

If you can climb one flight of stairs without getting chest pain or shortness of breath, you will probably be able to cope with the amount of energy required for sex. Avoid having sex after eating a large meal, drinking alcohol or when you are very tired.

You may find that some positions are more comfortable than others, because of discomfort in your chest wound. It can take time for sexual relationships to get back to normal after surgery, so try to take things slowly. Make an effort to discuss your feelings and concerns with your partner.



Quick tip

It's important that you are free of pain when you go to bed so that you get a good night's sleep. You might find it helpful to learn relaxation techniques or listen to relaxation tapes.

Quick tip

In the first few weeks after surgery, you might have trouble with your memory and concentration. This will improve steadily and should not be a problem.

Why am I so tired?

Tiredness is very common after surgery and it's important that you rest. Have a one to two hour rest in the afternoon. Even if you don't sleep, lie down and read a book or listen to relaxing music. Try to get eight to 10 hours of sleep every night.

Sometimes you may get breathless when you exercise. This is to be expected but, if it continues after you have rested, or is associated with heart 'flutters', tell your doctor. Breathlessness can be a symptom of heart attack (see page 12).

Why do I feel great one day, but dreadful the next?

'Up' and 'down' days after surgery are normal. The best way to overcome this is to plan something interesting to do each day. Don't alter your normal lifestyle too much either. Get out of bed at the usual time and dress in day clothes.

When can I return to work?

This decision is usually made after your four to six week check-up. It will depend on your type of work, how demanding your job is, your strength and other factors. Office workers can usually return to work in six weeks. If your work is more physically demanding, you should follow your surgeon's advice.

Palpitations

Many people have palpitations (thumping heartbeats or fluttery feelings) after operations. Usually this is not serious. If you are having palpitations often, or they last a long time, tell your doctor.


Heart attack

A heart attack is an emergency. Getting to hospital quickly can reduce the damage to your heart and increase your chance of survival.

A heart attack happens when there is a sudden, complete blockage of an artery that supplies blood to your heart muscle (a 'coronary artery'). As a result, some of your heart muscle starts to die. If the artery stays blocked, the lack of blood permanently damages the area of your heart muscle that is supplied by that artery.

It is important that you know what to do if you or someone else has a heart attack.

Will you recognise your heart attack?



Do you feel any

pain pressure heaviness tightness

In one or more of your

chest neck jaw arm/s back shoulder/s

You may also feel

nauseous a cold sweat dizzy short of breath

Yes

1 STOP and rest now

2 TALK Tell someone how you feel

If you take angina medicine

- Take a dose of your medicine.
- Wait 5 minutes. Still have symptoms? Take another dose of your medicine.
- Wait 5 minutes. Symptoms won't go away?

Are your symptoms severe or getting worse?

or

Have your symptoms lasted 10 minutes?

Yes

3 CALL 000* Triple Zero

- Ask for an ambulance.
- Don't hang up.
- Wait for the operator's instructions.

*If calling Triple Zero (000) does not work on your mobile phone, try 112.

© 2010 National Heart Foundation of Australia ABN 98 008 419 761.

Infection

Call your cardiologist or surgeon straight away if you have any symptoms of an infection, such as fever, sweating, loss of appetite, weight loss or an overall lack of energy lasting more than a week.

Endocarditis

People who have heart valve disease or who have had heart valve surgery are at greater risk than others of an infection called 'endocarditis'.

Endocarditis occurs when bacteria pass through the bloodstream and settle on the valve. This can cause very serious problems.

To prevent endocarditis, your cardiologist may tell you to take antibiotics before certain dental and surgical procedures and, in some cases, childbirth. If you take antibiotics, the risk of infection in these circumstances is very small. If you're not sure what antibiotics you should take or when you should take them, ask your doctor or surgeon.

Will I need to take medicine?

If you have had a valve replacement, you may need to take an anti-coagulant (anti-clotting) medicine. These medicines stop the blood clotting easily and are used to prevent clots forming on your new valve. Warfarin is the most commonly used anti-coagulant.

Some types of valve replacement require only two or four months of anti-coagulant treatment, but others require lifelong treatment. These medicines don't stop clotting completely, but reduce the risk of problems that can be caused by blood clots forming on your new valve.

Important things to remember about anti-coagulant medicine

- Don't stop taking anti-coagulants without talking to your doctor first.
- Don't change the amount of medicine that you are taking without talking to your doctor first.
- Try to take your medicine at the same time every day, preferably in the evening. If you miss a dose, take it as soon as you remember on the same day. If you don't remember until the next day, don't take an extra dose, but talk to your doctor.
- If you go away, make sure that you take enough medicine with you.
- To make sure that you take the right dose of medicine, you will need to have frequent blood tests to check the time it takes clots to form. While you are still in the hospital after your operation, your doctor or nurse will probably test your blood daily. When you go home at first, you should have a blood test once a week. Later, it may only need to be once every four to six weeks.
- While you are taking anti-coagulants, be on the alert for signs of bleeding and tell your doctor about any unusual bleeding or bruising.
- As other medicines can interfere with anti-coagulants, don't take any medicine that your doctor hasn't prescribed. This includes 'over the counter' medicines, such as cough medicines. For minor aches and pains, take paracetamol, not aspirin or ibuprofen (as these may affect clotting). Some people may be prescribed a regular dose of aspirin in addition to warfarin. If this is the case, it is important not to take extra aspirin for aches and pains.
- If you are taking warfarin, you should limit the amount of alcohol that you consume because it can alter the effectiveness of the medicine.
- If you need dental work or surgery, tell your dentist or surgeon that you are taking anti-coagulants. They may tell you to temporarily stop taking them.



- If you are planning a long trip, before you leave, ask your doctor to help you to make arrangements to have any necessary blood tests during your trip.
- Always carry with you information about your medicine and the dose that you are taking.
- Some foods can interfere with the way warfarin works in the body. See the box below for further information.

Warfarin and food

People taking warfarin need to be aware that certain types of foods can either enhance or inhibit the effects of the medicine. Foods high in vitamin K, such as leafy green vegetables, can reduce the effect of warfarin, while some dietary supplements may increase the effects of warfarin.

It is important to remember that these foods should not be avoided altogether, but that people taking warfarin should try to keep their intake of these foods fairly consistent.

Foods to be aware of include:

- leafy green vegetables (such as spinach, broccoli, lettuce, cabbage, asparagus, parsley, watercress and brussels sprouts)
- canola, olive and soybean oils
- seaweed (such as in sushi)
- green and herbal teas
- liver
- spring onions.

People taking warfarin should talk to a doctor first if they are considering changes to their eating habits (for example, switching to a vegetarian diet, following a special meal plan to lose or gain weight, or starting or discontinuing any dietary supplements, such as fish oil supplements).

For more information, contact the NPS Medicines Line on 1300 633 424.

What should I talk to my doctor about?

Heart surgery is a major event in your life and you should make sure that you are well informed about why you need surgery, and what the benefits and risks of surgery are. Below are some questions about heart valve surgery that you may want to ask your doctor.

- What benefits will I get from the surgery?
- What are the risks or potential complications of surgery?
- Will I need a blood transfusion and can I donate blood for myself?
- What sort of heart valve will I receive?
- Will I need to take anti-coagulant medicine? If so, for how long?
- How much will the operation cost and how much will be covered by insurance?
- Is there a place nearby where my partner can stay while I'm in hospital?
- If I need to travel from my home to a different town for the operation, how long should I stay there after my discharge?
- How often should I have follow-up visits?
- What are the symptoms of heart valve failure?

Glossary

Annuloplasty rings

Special rings used to narrow an enlarged valve and make repairs stronger.

Anti-coagulants

Medicines used to treat and prevent blood clotting inside the blood stream.

Aorta

The main artery of the body. The aorta rises directly from the left ventricle (the main pumping chamber of the heart) and supplies oxygen-rich blood to all other arteries except the pulmonary artery.

Aortic valve

The valve separating the left ventricle and the aorta.

Atrium (plural = atria)

One of the heart's two upper (collecting) chambers.

Bacterial endocarditis

An infection of the heart valves.

Balloon valvotomy

A procedure using a balloon to open stuck valves.

Biological tissue valves

Valves made from human or animal tissue.

Heart murmur

A swishing sound caused by blood flowing forwards or backwards in the heart abnormally.

Incompetent valve

A 'leaking' valve that lets blood to flow back into a chamber of the heart.

Mechanical valves

Valves made from non-biological materials, such as plastic or metal.

Mitral valve

The valve between the left atrium and left ventricle.

Murmur

See 'heart murmur'.

Prosthetic rings

See 'annuloplasty rings'.

Pulmonary valve

The valve separating the right ventricle and the pulmonary artery.

Regurgitant valve

See 'incompetent valve'.

Stenotic valve

A narrowed, stiff valve.

Tricuspid valve


The valve between the right atrium and right ventricle.

Ventricle

One of the two main pumping chambers of the heart.

Warfarin

One of a group of anti-coagulant (anti-clotting) medicines.



For heart health information
1300 36 27 87
www.heartfoundation.org.au

Key things to remember about heart valve surgery

Heart valves are like one-way doors that control the direction of blood flow between the four chambers of the heart.

Sometimes people are born with damaged valves, but disease or infection can also damage your valves.

There are two main types of heart valve surgery that you can have to fix a damaged or faulty heart: valve repair and valve replacement.

Normal recovery time from heart surgery takes four to six weeks. During this time, you will begin to strengthen your muscles and return to your usual activities.

If you have had a valve replacement, you may need to take an anti-coagulant (anti-clotting) medicine, such as warfarin.

The Heart Foundation and World Health Organization recommend that all patients who have had a heart attack, heart surgery, coronary angioplasty, angina or other heart and blood vessel disease attend an appropriate cardiac rehabilitation and prevention program.

© 2015 National Heart Foundation of Australia ABN 98 008 419 761

ISBN 978-1-921748-65-3

First published 2010. Reprinted with corrections 2011, 2013, 2015. CON-083.v4

Terms of use: This material has been developed for general information and educational purposes only. It does not constitute medical advice. Please consult your healthcare provider if you have, or suspect you have, a health problem. The health information provided has been developed by the Heart Foundation and is based on independent research and the available scientific evidence at the time of writing. The information is obtained and developed from a variety of sources including, but not limited to, collaborations with third parties and information provided by third parties under licence. It is not an endorsement of any organisation, product or service.

While care has been taken in preparing the content of this material, the National Heart Foundation of Australia, its employees and related parties cannot accept any liability, including for any loss or damage, resulting from the reliance on the content, or for its accuracy, currency and completeness.

This material may be found in third parties' programs or materials (including, but not limited to, show bags or advertising kits). This does not imply an endorsement or recommendation by the National Heart Foundation of Australia for such third parties' organisations, products or services, including their materials or information. Any use of National Heart Foundation of Australia materials or information by another person or organisation is at the user's own risk.

The entire contents of this material are subject to copyright protection. Enquiries concerning permissions should be directed to copyright@heartfoundation.org.au